

## Last time: Harmonic Functions

The Laplacian operator, written  $\nabla^2$  or  $\Delta$ , acts on functions  $g : \mathbb{R}^2 \rightarrow \mathbb{R}$  by

$$\nabla^2 g = \nabla \cdot \nabla g = \frac{\partial^2 g}{\partial x^2} + \frac{\partial^2 g}{\partial y^2}$$

### Definition

A function  $u : \mathbb{R}^2 \rightarrow \mathbb{R}$  is *harmonic* if  $\nabla^2 u = 0$ .

### Lemma

Let  $f(z) = u(x, y) + iv(x, y)$  be analytic on a domain  $D$ . Then  $u$  and  $v$  are harmonic on  $D$ .

### Can you go backwards?

Yes, if  $u$  defined on a *simply-connected* region.

# When is $u$ the real part of analytic functions?

From the 2012-2013 exam

(iii) Find all the functions  $k$  analytic on  $\mathbb{C}$  with  $\operatorname{Re}(k(x+iy)) = 2x - \sinh x \sin y$ , giving an explicit expression for  $k(z)$  in terms of  $z$ . Show that you have found **all** the functions satisfying the above conditions. *(6 marks)*

The real part of an analytic function is harmonic

First, check  $\nabla^2 u = 0$ . If not, the answer is no.

If it is, find  $f'$  using Cauchy-Riemann

$$f' = \frac{\partial}{\partial x} (u(x, y) + iv(x, y)) = u_x + iv_x = u_x - iu_y$$

This gives us  $f'$  in terms of  $x$  and  $y$ . We'd *like* to write  $f'$  in terms of  $z$ , and integrate to find  $f$ . But how?

Maybe we need a clever little trick....

## Dr. Hart's "Clever little trick"

Given:

We know  $f'$  in terms of  $x$  and  $y$ , want it terms of  $z$ .

Guess:

Set  $y = 0$ ; to get  $f'(x)$  in terms of just  $x$ . Integrate to get  $f(x)$ .

Guess that this is actually formula for  $f(z)$

Check:

Show that  $\operatorname{Re}(f) = u(x, y)$

To find *all* such  $k$ :

Lemma

*Suppose that  $f$  and  $g$  are analytic on a region  $D$  and that  $\operatorname{Re}(f) = \operatorname{Re}(g)$  on  $D$ . Then  $f = g + ia$  for some  $a \in \mathbb{R}$ .*

## A last application of Cauchy-Riemann

- ▶ Cauchy-Riemann relates between  $\operatorname{Re}(f)$  and  $\operatorname{Im}(f)$
- ▶ If we have more relations, then  $f$  is *very* constrained

### Example from the notes:

The function  $f$  is analytic in  $\mathbb{C}$  and its real and imaginary parts  $u$  and  $v$  satisfy

$$ue^v = 12$$

at all points in  $\mathbb{C}$ . Prove that  $f$  is constant.

### Next up – Section 6: Power series

- ▶ Not much different than real power series, so largely review
- ▶ Spoiler: analytic functions have convergent power series